



A Reliable Research Partner in Life Science and Medicine

Elab Fluor® 488 Anti-Human CD161 Antibody[HP-3G10]

Catalog Number: E-AB-F1155L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| _ | | | | | |
|---|-----|-------|---|-----|-----|
| | esc | - IVI | | 17 | n n |
| ш | しつし | 71 II | ы | 41. | 9 |

Reactivity Human Host Mouse

Isotype Mouse IgG1, κ
Clone No. HP-3G10

Isotype Control Elab Fluor® 488 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792L]

Conjugation Elab Fluor® 488

Conjugation Information Elab Fluor® 488 is designed to be excited by the Blue laser (488 nm) and detected using

an optical filter centered near 520 nm (e.g., a 525/40 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names NKRP1A;CLEC5B;HNKR-P1a;KLRB1;NKR-P1A

 Uniprot ID
 Q12918

 Gene ID
 3820

Background CD161 is a type II transmembrane glycoprotein, also known as NKR-P1A, that is

expressed as a 40-44 kD homodimer. It is a member of the C-type lectin superfamily. CD161 is expressed on a majority of NK cells, NKT cells, and subsets of peripheral T cells and CD3+ thymocytes. It has been reported that Th17 cells are a subpopulation of CD4+CD161+CCR6+ cells. While the biological function of CD161 is not clear, it has been suggested to serve either as a stimulatory receptor or to inhibit NK cell-mediated cytotoxicity and cytokine production. LLT-1 (lectin-like transcript-1, also named as

osteoclast inhibitory lectin or CLEC2D) is the ligand of CD161.

Web: www.elabscience.cn

For Research Use Only

Rev. V1.5