Recombinant Human SIGLEC2/CD22 Protein (His Tag)

Catalog Number: PDMH100369

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Species	Human		
Source	HEK293 Cells-derived Human SIGLEC2 protein Asp20-Arg687, with an C-terminal His		
Calculated MW	76.2 kDa		
Observed MW	100-140 kDa		
Accession	P20273		
Bio-activity	Not validated for activity		
Properties			
Purity	> 95% as determined by reducing SDS-PAGE.		
Endotoxin	< 1.0 EU/mg of the protein as determined by the LAL method		
Storage	ge Generally, lyophilized proteins are stable for up to 12 months when stored at -20		
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of		
	reconstituted samples are stable at $< -20^{\circ}C$ for 3 months.		
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.		
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with 5% Trehalose and 5%		
	Mannitol.		
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of		
	0.5 mg/mL. Concentration is measured by UV-Vis.		

Data

KDa	М	R
80 60	-	-
60		
40	-	
30	-	
20	-	
12	-	

> 95 % as determined by reducing SDS-PAGE.

Background

Mediates B-cell B-cell interactions. May be involved in the localization of B-cells in lymphoid tissues. Binds sialylated glycoproteins, one of which is CD45. Preferentially binds to alpha-2,6-linked sialic acid. The sialic acid recognition site can be masked by cis interactions with sialic acids on the same cell surface. Upon ligand induced tyrosine phosphorylation in the immune response seems to be involved in regulation of B-cell antigen receptor signaling. Plays a role in positive regulation through interaction with Src family tyrosine kinases and may also act as an inhibitory receptor by recruiting cytoplasmic phosphatases via their SH2 domains that block signal transduction through dephosphorylation of signaling molecules.

Web:www.elabscience.com