Recombinant Mouse IL-2RG/CD132 Protein (His Tag)

Catalog Number: PKSM041317

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Species			Mouse
Source			HEK293 Cells-derived Mouse IL-2RG/CD132 protein Ser25-Ala263, with an C-terminal
			His
Calculated MW			29 kDa
Observed MW			45-75 kDa
Accession			Q3UPA9
Bio-activity			Not validated for activity
Properties			
Purity			> 95 % as determined by reducing SDS-PAGE.
Endotoxin			< 1.0 EU per µg of the protein as determined by the LAL method.
Storage			Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
			°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
			reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
Shipping			This product is provided as lyophilized powder which is shipped with ice packs.
Formulation			Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.5.
			Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
			before lyophilization.
			Please refer to the specific buffer information in the printed manual.
Reconstitution			Please refer to the printed manual for detailed information.
Data			
	kDa	МК	R
	120 90	the state	
	90 60	and a	
	40	Torrest	
	70	1	
	30	1200	
	20		

Background

The Interleukin-2 receptor gamma chain (IL-2 R γ , CD132) of the high affinity functional mouse IL-2 receptor complex is a member the hematopoietin receptor family. It is expressed on most lymphocyte (white blood cell) populations, and its gene is found on the X-chromosome of mammals. Common IL2 receptor- γ Chain is required for IL-2 receptor signaling. Besides IL-2, the Common IL2 receptor- γ chain has been shown to be a component of the functional receptor complexes for IL-4, IL-7, IL-9 and IL-15. It is a component of multiple cytokine receptors that are essential for lymphocyte development and function. Common IL2 receptor- γ Chain is also designated the common gamma chain.

14