Elabscience Biotechnology Co., Ltd.



A Reliable Research Partner in Life Science and Medicine

Mouse ST2 Antibody Pair Set

Catalog No.E-KAB-0606ApplicationsELISASynonymsDER4;FIT-1;Growth stimulation expressed gene 2;IL-1R4;IL1RL1;IL33R;ST2

Kit components & Storage

Title	Specifications	Storage
Mouse ST2 Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze/thaw cycles.
Mouse ST2 Detection Antibody (Biotin)	1 vial, 50 μL	Store at -20°C for one year.
		Avoid freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0606)		
		Mouse ST2 Capture Antibody	Mouse ST2 Detection Antibody	
			(Biotin)	
Immunogen	Immunogen	Recombinant Mouse ST2 protien	Recombinant Mouse ST2 protien	
Information	Swissprot	P14719		
Product details	Reactivity	Mouse	Mouse	
	Host	Goat	Goat	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5 mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%	
		glycerol; pH 7.5	protective protein; 50% glycerol; pH	
			7.5	
	Purify	Antigen Affinity	Antigen Affinity	
	Specificity	Detects Mouse ST2 in ELISAs.		

Tel: 400-999-2100 Web: www.elabscience.cn Email: techsupport@elabscience.cn





A Reliable Research Partner in Life Science and Medicine

Applications

Mouse ST2 Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Mouse ST2 Capture	
Capture		Antibody	10
			Attention
ELISA	1:1000-1:10000	Mouse ST2 Detection	Optical Density
Detection		Antibody (Biotin)	0,1
			•
			0.1 10 100 Mouse ST2 Concentration (ng/mL)
			mouse or 2 concentration (fightle)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

The protein encoded by this gene is a member of the interleukin 1 receptor family. Studies of the similar gene in mouse suggested that this receptor can be induced by proinflammatory stimuli , and may be involved in the function of helper T cells. This gene , interleukin 1 receptor , type I (IL1R1) , interleukin 1 receptor , type II (IL1R2) and interleukin 1 receptor-like 2 (IL1RL2) form a cytokine receptor gene cluster in a region mapped to chromosome 2q12. Alternative splicing of this gene results in multiple transcript variants.

Tel: 400-999-2100 Web: www.elabscience.cn Email: techsupport@elabscience.cn