

A Reliable Research Partner in Life Science and Medicine

Elab Fluor® 488 Anti-Human CD8a Antibody[OKT-8]

Catalog Number: E-AB-F1110L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human Mouse Host

Isotype Mouse IgG2a, ĸ

OKT-8 Clone No.

Isotype Control Elab Fluor[®] 488 Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802L]

Conjugation Elab Fluor®488

Conjugation Information Elab Fluor® 488 is designed to be excited by the Blue laser (488 nm) and detected using

an optical filter centered near 520 nm (e.g., a 525/40 nm bandpass filter).


Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer. Storage Buffer

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Human peripheral blood lymphocytes are stained with Elab

Fluor® 488 Anti-Human CD8a Antibody (filled gray histogram). Unstained lymphocytes (empty black histogram) are used as control.

Preparation & Storage

Keep as concentrated solution. **Storage**

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD8A;MAL:T-cell surface glycoprotein CD8 alpha chain:T-lymphocyte differentiation

antigen T8/Leu-2

Uniprot ID P01732 Gene ID 925

For Research Use Only

Fax: 1-832-243-6017 Tel: 1-832-243-6086 Toll-free: 1-888-852-8623 Email:techsupport@elabscience.com

Web:www.elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Background

CD8a is a 32-34 kD type I glycoprotein. It forms a homodimer (CD8a/a) or heterodimer (CD8a/b) with CD8b. CD8, also known as T8 and Leu2, is a member of the immunoglobulin superfamily found on the majority of thymocytes, a subset of peripheral blood T cells, and NK cells (which express almost exclusively CD8a homodimers). CD8 acts as a co-receptor with MHC class I-restricted T cell receptors in antigen recognition and T cell activation and has been shown to play a role in thymic differentiation. Two domains in CD8a are important for function: the extracellular IgSF domain binds the a3 domain of MHC class I and the cytoplasmic CXCP motif binds the tyrosine kinase p56 Lck.

Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web:www.elabscience.com