

HRP-conjugated PCNA Monoclonal Antibody

catalog number: AN00472HP

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human;Mouse;Rat
Immunogen	Recombinant human PCNA protein expressed by E.coli
Host	Mouse
Isotype	IgG2a
Clone	9C6
Purification	Protein A/G Purification
Conjugation	HRP
Buffer	PBS with 0.05% Proclin300, 1% protective protein and 50% glycerol, pH7.4

Applications

Recommended Dilution

WB	1:2500-1:5000
----	---------------

Data



Western blot with HRP-conjugated PCNA Monoclonal Antibody at dilution of 1:5000. lane 1: Raji whole cell lysate, lane 2: HepG2 whole cell lysate, lane 3: HL-60 whole cell lysate, lane 4: Raw264.7 whole cell lysate, lane 5: NIH/3T3 whole cell lysate, lane 6: PC-12 whole cell lysate

Observed-MW:29 kDa

Calculated-MW:29 kDa

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Protected from prolonged exposure to light.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

Auxiliary protein of DNA polymerase delta and epsilon, is involved in the control of eukaryotic DNA replication by increasing the polymerase's processibility during elongation of the leading strand. Induces a robust stimulatory effect on the 3'-5' exonuclease and 3'-phosphodiesterase, but not apurinic-apyrimidinic (AP) endonuclease, APEX2 activities. Has to be loaded onto DNA in order to be able to stimulate APEX2. Plays a key role in DNA damage response (DDR) by being conveniently positioned at the replication fork to coordinate DNA replication with DNA repair and DNA damage tolerance pathways. Acts as a loading platform to recruit DDR proteins that allow completion of DNA replication after DNA damage and promote postreplication repair: Monoubiquitinated PCNA leads to recruitment of translesion (TLS) polymerases, while 'Lys-63'-linked polyubiquitination of PCNA is involved in error-free pathway and employs recombination mechanisms to synthesize across the lesion.