

Monkey GDF15 Antibody Pair Set

Catalog No.	E-KAB-0669	Applications	ELISA
Synonyms	GDF-15;MIC-1;MIC1;NAG-1;PDF;PLAB;PTGFB;TGF-PL		

Kit components & Storage

Title	Specifications	Storage
Monkey GDF15 Capture Antibody	1 vial, 100 µg	Store at -20°C for one year. Avoid freeze/thaw cycles.
Monkey GDF15 Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20°C for one year. Avoid freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Product Information

Items		Characteristic (E-KAB-0669)	
		Monkey GDF15 Capture Antibody	Monkey GDF15 Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Monkey GDF15 protein	Recombinant Monkey GDF15 protein
	Swissprot	G7PWZ3	
Product details	Reactivity	Monkey	Monkey
	Host	Goat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300; 50% glycerol; pH 7.5	PBS with 0.04% Proclin 300; 1% protective protein; 50% glycerol; pH 7.5
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Monkey GDF15 in ELISAs.	

For Research Use Only

Applications

Monkey GDF15 Sandwich ELISA Assay

	Recommended Concentration/Dilution	Reagent	Images																						
ELISA Capture	0.5-4 µg/mL	Monkey GDF15 Capture Antibody	 <p>The graph is a standard curve for the ELISA assay. The x-axis represents Monkey GDF15 Concentration in pg/mL on a logarithmic scale from 1 to 10,000. The y-axis represents Optical Density on a logarithmic scale from 0.01 to 10. The data points show a clear upward trend, indicating that as the concentration of Monkey GDF15 increases, the optical density also increases.</p> <table border="1"> <caption>Approximate data points from the standard curve</caption> <thead> <tr> <th>Monkey GDF15 Concentration (pg/mL)</th> <th>Optical Density</th> </tr> </thead> <tbody> <tr><td>10</td><td>0.05</td></tr> <tr><td>20</td><td>0.1</td></tr> <tr><td>50</td><td>0.2</td></tr> <tr><td>100</td><td>0.4</td></tr> <tr><td>200</td><td>0.8</td></tr> <tr><td>500</td><td>1.5</td></tr> <tr><td>1000</td><td>2.5</td></tr> <tr><td>2000</td><td>4.0</td></tr> <tr><td>5000</td><td>6.0</td></tr> <tr><td>10000</td><td>8.0</td></tr> </tbody> </table>	Monkey GDF15 Concentration (pg/mL)	Optical Density	10	0.05	20	0.1	50	0.2	100	0.4	200	0.8	500	1.5	1000	2.5	2000	4.0	5000	6.0	10000	8.0
Monkey GDF15 Concentration (pg/mL)	Optical Density																								
10	0.05																								
20	0.1																								
50	0.2																								
100	0.4																								
200	0.8																								
500	1.5																								
1000	2.5																								
2000	4.0																								
5000	6.0																								
10000	8.0																								
ELISA Detection	1:1000-1:10000	Monkey GDF15 Detection Antibody (Biotin)																							

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Bone morphogenetic proteins (e.g. , BMP9 , MIM 605120) are members of the transforming growth factor-beta (see TGFB1 , MIM 190180) superfamily and regulate tissue differentiation and maintenance. They are synthesized as precursor molecules that are processed at a dibasic cleavage site to release C-terminal domains containing a characteristic motif of 7 conserved cysteines in the mature protein. GDF15 mRNA is most abundant in the liver , with lower levels seen in some other tissues. Its expression in liver can be significantly up-regulated in during injury of organs such as liver , kidney , heart and lung.