## **Elabscience**®

### **SEPT8 Polyclonal Antibody**


### catalog number: E-AB-65176

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of human 44082 (NP_001287728.1).
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution


WB	1:500-1:2000
IF	1:50-1:200

#### Data




Western blot analysis of extracts of various cell lines using SEPT8 Polyclonal Antibody at dilution of 1:1000.

Observed-MW:54 kDa Calculated-MW:43 kDa/49 kDa/51 kDa/55 kDa



Immunofluorescence analysis of C6 cells using SEPT8 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of L929 cells using SEPT8 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining. Immunofluorescence analysis of U-2 OS cells using SEPT8 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

#### Background

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w .elabscience.com</u>

Tel: 1-832-243-6086 Email:techsupport@elabscience.com

# **Elabscience**®

This gene is a member of the septin family of nucleotide binding proteins, originally described in yeast as cell division cycle regulatory proteins. Septins are highly conserved in yeast, Drosophila, and mouse, and appear to regulate cytoskeletal organization. Disruption of septin function disturbs cytokinesis and results in large multinucleate or polyploid cells. Multiple alternatively spliced transcript variants encoding different isoforms have been found for this gene.

For Research Use Only