A Reliable Research Partner in Life Science and Medicine

AF/LE Purified Anti-Mouse CD106 Antibody[M/K-2.7]

catalog number: E-AB-F10910

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity	Mouse
Host	Rat

Isotype Rat IgGl, κ **Clone** M/K-2.7


Buffer Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method

.

Applications	Recommended	Dilution
--------------	-------------	----------

FCM $2 \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$

Data



C57/BL6 Mouse bone marrow cells were stained with 0.2 µg

AF/LE Purified Anti-Mouse CD106 Antibody[M/K-2.7]

(Right) and 0.2 μg Rat IgG1, κ Isotype Control (Left),

followed by PE-conjugated Goat Anti-Rat IgG Secondary

Antibody.

Preparation & Storage

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles. This preparation contains no preservatives, thus it should be handled

under aseptic conditions.

Shipping Ice bag

Background

CD106 is a 110 kD glycosylphosphatidylinositol (GPI)-linked transmembrane protein, also known as VCAM-1 and INCAM-110. It is constitutively expressed on bone marrow stromal cells, myeloid progenitors, splenic dendritic cells, activated endothelial cells, as well as some lymphocytes. CD106 expression can be upregulated on endothelial cells by inflammatory cytokines. CD106 is involved in adhesion and acts as a counter-receptor for VLA-4 (α 4/ β 1 integrin) and LPAM-1 (α 4/ β 7 integrin).

For Research Use Only

Fax: 1-832-243-6017