

CCL15/MIP-1 δ Polyclonal Antibody(Capture/Detector)

catalog number: AN004500P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Immunogen	Recombinant Human CCL15/MIP-1 δ Protein expressed by E.coli
Host	Rabbit
Isotype	Rabbit IgG
Purification	Antigen Affinity Purification
Conjugation	Unconjugated
buffer	Phosphate buffered solution, pH 7.2, containing 0.05% proclin 300.

Applications

Recommended Dilution

ELISA Capture	2-8 μ g/mL
ELISA Detector	0.1-0.4 μ g/mL

Data



Sandwich ELISA-Recombinant Human CCL15/MIP-1 δ Protein standard curve. Background subtracted standard curve using Anti-CCL15/MIP-1 δ antibody(AN004500P) (Capture), Anti-CCL15/MIP-1 δ antibody(AN004500P) (Detector). The reference range value is 15.6 - 1,000 pg/mL for human.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

Chemokine ligand 15 (CCL15), also known as leukotactin-1, MIP5, MIP1 and HCC-2, is a small cytokine belonging to the C-C chemokine family. CCL15 is prevalently expressed in liver, small intestine, colon, and in certain leukocytes and macrophages of the lung. It is chemotactic for neutrophils, monocytes, and lymphocytes and elicits its effects by binding to cell surface chemokine receptors like CCR1 and CCR3.

For Research Use Only