

Recombinant Human PDGFRα/CD140a Protein (His Tag)

Catalog Number: PKSH033564

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human PDGFRα/CD140a protein Gln24-Glu524, with an C-terminal His
Calculated MW	57.0 kDa
Observed MW	93 kDa
Accession	P16234
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Platelet-derived Growth Factor Receptor Alpha (PDGF Rα) is an enzyme that belongs to the class III subfamily of receptor tyrosine kinases. It is a type I transmembrane glycoprotein, and can form homo- or hetero-dimeric receptors when engaged by dimers of the PDGF family of growth factors. PDGF Rα is strongly expressed in oligodendrocyte, lung, skin and intestinal progenitor cells and induced by inflammation or growth in culture, but is lowly expressed in most mesenchymal cells. PDGF Rα autophosphorylates upon dimerization, activating signaling cascades in PI-3kinase Ras-MAP kinase, and PLC-γ pathways. PDGF Rα has influence on local gradients of epithelially produced PDGF-AA or PDGF-CC during formation of the cranial, cardiac neural crest and interstitial kidney mesenchyme.

For Research Use Only