

# Recombinant Human PIK3CA Protein (His Tag)

Catalog Number:PDEH100021



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## Description

<b>Synonyms</b>	5-bisphosphate 3-kinase 110 kDa catalytic subunit alpha;5-bisphosphate 3-kinase catalytic subunit alpha isoform;caPI3K;CLOVE;CWS5;MCAP;MCM;MCMTC;MGC142161;MGC142163;p110 alpha;p110alpha;Phosphatidylinositol 3 kinase catalytic alpha polypeptide;Phosphatidylinositol 3 kinase catalytic 110 KD alpha;Phosphatidylinositol 4 5 bisphosphate 3 kinase catalytic subunit alpha;Phosphatidylinositol 4 5 bisphosphate 3 kinase catalytic subunit alpha isoform;Phosphatidylinositol 4;5 bisphosphate 3 kinase 110 kDa catalytic subunit alpha;Phosphatidylinositol-4;Phosphoinositide 3 kinase catalytic alpha polypeptide;PI3 kinase p110 subunit alpha;PI3-kinase subunit alpha;PI3K;PI3K-alpha;PI3KC A;PIK3C A;Pik3ca;PK3CA;PK3CA;PtdIns 3 kinase p110;PtdIns-3-kinase subunit alpha;PtdIns-3-kinase subunit p110-alpha;Serine/threonine protein kinase PIK3CA
<b>Species</b>	Human
<b>Expression Host</b>	E.coli
<b>Sequence</b>	Met1-Pro200
<b>Accession</b>	P42336-1
<b>Calculated Molecular Weight</b>	23.1 kDa
<b>Observed molecular weight</b>	25.4 kDa
<b>Tag</b>	N-His

## Properties

<b>Purity</b>	> 90 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Please contact us for more information.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Background

Phosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a 110 kDa catalytic subunit. The protein encoded by this gene represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has been found to be oncogenic and has been implicated in cervical cancers. A pseudogene of this gene has been defined on chromosome 22.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: [www.elabscience.com](http://www.elabscience.com)

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)