Recombinant Mouse IL-17F protein(His Tag)

Catalog Number: PKSM041467



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| - | | | |
|--------------|-----|-----|------|
| 1116 | 100 | rin | ion |
| \mathbf{L} | | ттр | tion |

 Species
 Mouse

 Mol_Mass
 15.8 kDa

 Accession
 Q7TNI7

Bio-activity Measure by its ability to induce IL-6 secretion in 3T3 cells. The ED₅₀ for this effect is

<100 ng/mL.

Properties

Purity > 98 % as determined by reducing SDS-PAGE.

Endotoxin < 0.1 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile 20 mM sodium citrate, pH 4.5.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

Interleukin-17 is a potent pro-inflammatory cytokine produced by activated memory T cells. There are at least six members of the IL-17 family in humans and in mice. Today; IL-17 represents a family of structurally related cytokines that share a highly conserved C-terminal region but differ from one another in their N-terminal regions and in their distinct biological roles. The six known members of this family; IL-17A through IL-17F; are secreted as homodimers. IL-17F also regulates cartilage matrix turnover and inhibits angiogenesis.

For Research Use Only