

# Recombinant Mouse SIRPA/CD172a Protein (Fc Tag)

Catalog Number:PKSM041145



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	Tyrosine-Protein Phosphatase Non-Receptor Type Substrate 1;CD172 Antigen-Like Family Member A;Signal-Regulatory Protein Alpha-1;CD172a;SIRPA;MFR;SHPS1;SIRP
<b>Species</b>	Mouse
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Lys32-Asn373
<b>Accession</b>	Q6P6I8
<b>Calculated Molecular Weight</b>	64.8 kDa
<b>Observed molecular weight</b>	115-125 kDa
<b>Tag</b>	C-Fc

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Mouse Signal Regulatory Protein  $\alpha$  (SIRP $\alpha$ ) is a type I transmembrane glycoprotein. It contains two Ig-like C1-type domains and one Ig-like V-type domain. Mouse SIRP alpha ECD shares 61%, 75%, 62%, 61%, and 59% aa sequence identity with human, rat, equine, bovine, and porcine SIRP alpha, respectively. SIRP $\alpha$  can express in various tissues, mainly on brain and myeloid cells, including macrophages, neutrophils, dendritic and Langerhans cells. It also can detect in neurons, smooth muscle and endothelial cells. SIRPA is an immunoglobulin-like cell surface receptor for CD47. SIRP $\alpha$  acts as docking protein and induces translocation of PTPN6, PTPN11 and other binding partners from the cytosol to the

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Mouse SIRPA/CD172a Protein (Fc Tag)

Catalog Number:PKSM041145



plasma membrane. SIRP $\alpha$  shows adhesion of cerebellar neurons, neurite outgrowth and glial cell attachment. SIRP $\alpha$  engagement generally produces a negative regulatory signal; it may mediate negative regulation of phagocytosis, mast cell activation and dendritic cell activation

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: [www.elabscience.com](http://www.elabscience.com)

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)