

Anti-Human CD3-FITC/CD16+CD56-PE Cocktail

Catalog Number: E-AB-FC0001

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Clone No.	OKT-3;3G8;5.1H11
Conjugation	FITC;PE
Conjugation Information	FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical filter centered near 530 nm (e.g., a 525/40 nm bandpass filter). PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green (561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications

Recommended usage

FCM	For whole blood samples, add 5 µL Anti-Human CD3-FITC/CD16+CD56-PE Cocktail to 100 µL anticoagulant-treated blood sample. Mix and incubate the sample at 4°C in the dark for 30 min. Remove red blood cells with RBC lysis solution following the manufacturer's instruction. Wash the cell with cell staining buffer and discard the supernatant after centrifugation at 300 g for 5 min. Resuspend the cells with 200 µL cell staining buffer and load the sample on flow cytometer for detection. For other samples, 1×10^6 dissociated single cells are centrifuged at 300 g for 5 min with the supernatant discarded. Resuspend the cells with 100 µL cell staining buffer and add 5 µL Anti-Human CD3-FITC/CD16+CD56-PE Cocktail. Mix and incubate the sample at 4°C in the dark for 30 min. Add cell staining buffer to each tube, centrifuge at 300 g for 5 min and discard the supernatant. Resuspend the cells with 200 µL cell staining buffer and load the sample on flow cytometer for detection.
------------	---

Data



Human peripheral blood lymphocytes are stained with Anti-Human CD3-FITC/CD16+CD56-PE.

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

For Research Use Only

Background

This product is a FCM antibody cocktail made up of FITC Anti-Human CD3 Antibody [Clone: OKT-3] (Mouse IgG2a, κ), PE Anti-Human CD16 Antibody [Clone: 3G8] (Mouse IgG1, κ) and PE Anti-Human CD56 Antibody [Clone: 5.1H11] (Mouse IgG1, κ).; CD3 is a heterotetrameric protein consisting of a CD3 γ , a CD3 δ and 2 CD3 ϵ . It forms complex with TCR. OKT-3 recognize human CD3 ϵ . Human CD3 is expressed on the surface of T cells and NKT cells.;

CD16 is a low affinity receptor for the Fc of IgG. Human CD16 has two isoforms, CD16a and CD16b. CD16a is expressed on NK cells, activated monocytes and macrophages. CD16b is expressed on neutrophils. NK cells exert the function of ADCC by binding Fc of IgG through CD16.;

CD56 is also call neural cell adhesion molecule (NCAM), expressed on neurons, glia and skeletal muscle cells. In hematopoietic cells, CD56 is also expressed on NK cells and NKT cells. CD56 can be used to detect NK cells, $\gamma\delta$ T cells and activated CD8+ cells.