

## Recombinant Mouse SIGIRR/TIR8 Protein (His &Fc Tag)

**Catalog Number:** PKSM040853

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Mouse
<b>Source</b>	HEK293 Cells-derived Mouse SIGIRR/TIR8 protein Met 1-His 117, with an C-terminal His & Fc
<b>Calculated MW</b>	40.7 kDa
<b>Observed MW</b>	60-65 kDa
<b>Accession</b>	NP_075546.2
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

#### For Research Use Only

Single Ig IL-1-related receptor (SIGIRR) or TIR8 is a member of Toll-like receptor-interleukin 1 receptor signaling (TLR-IL-1R) receptor superfamily. Although SIGIRR/TIR8 shows the typical conserved motifs that characterize the IL-1R and Toll superfamily, it is structurally and functionally distinct from both. SIGIRR/TIR8 has only one Ig domain in its extracellular portion whereas the IL-1R family contains three Ig folds. An unusually long cytoplasmic domain is reminiscent of the structure of drosophila Toll, yet the SIGIRR peptide sequence is more closely related to IL-1RI. SIGIRR/TIR8 was mainly expressed in mouse and human epithelial tissues such as kidney, lung and gut. Resting and activated T and B lymphocytes and monocytes-macrophages expressed little or no SIGIRR/TIR8, with the exception of the mouse GG2EE macrophage line. Inflammation is enhanced in SIGIRR-deficient mice. SIGIRR negatively modulates immune responses. Inflammation is enhanced in SIGIRR-deficient mice, as shown by their enhanced chemokine induction after IL-1 injection and reduced threshold for lethal endotoxin challenge.

## For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017