Recombinant Human CD7/GP40 Protein (Fc Tag)

Catalog Number: PKSH033512



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-						٠,		
D	Α		C	PТ	m	ŤΤ	U.	m
\mathbf{L}	v	0	v.	ш	עו	UΙ	U.	

 Species
 Human

 Mol_Mass
 43.4 kDa

 Accession
 P09564

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

T-Cell Antigen CD7 is a single-pass type I membrane protein that that belongs to the the immunoglobulin superfamily. Human CD7 is synthesized as a 240 amino acid precursor that contains a 25 amino acid signal sequence and a 215 amino acid mature chain with a Ig-like (immunoglobulin-like) domain. CD7 is normally expressed on all T-lymphocytes; NK-cell s; pre-B lymphocytes and pleuripotent hematopoietic stem cells. CD7 plays an essential role in T-cell interactions; T-cell/B-cell interaction during early lymphoid development; T- and NK-cell activation and cytokine production. CD7 has been shown to interact with PIK3R1and SECTM1. However; the function of the CD7 protein in the immune system is still largely unknown.

For Research Use Only