# Recombinant Human Activin Receptor 2B/ACVR2B Protein (Fc &His Tag)

Catalog Number: PKSH032041



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-						
	AC	CI	чTГ	ΥП	on	
$\mathbf{L}$	$\mathbf{c}$	UI	ш	JUL	$\mathbf{v}_{\mathbf{H}}$	

 Species
 Human

 Mol\_Mass
 41.3 kDa

 Accession
 O13705

**Bio-activity** Not validated for activity

### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Activin proteins that belong to the transforming growth factor-beta (TGF-β) superfamily; exert their biological actions by binding to heteromeric receptor complexes of type I and type II serine/threonine kinase receptors. On ligand binding; type I and II receptors form a stable complex; resulting in phosphorylation of type I receptors by type II receptors with constitutive kinase activity; and subsequently initiates the activation of downstream molecules including the endogenous Smads. ActRIIB; also known as ActRIIB; is a type II receptor containing an extracellular domain (ECD); a transmembrane segment; and a cytoplasmic region that includes the kinase domain. ActRIIB is a receptor for activin A; activin B and inhibin A. Multiple ActRIIB isoforms can also be generated; which bind activin isoforms with different affinities.

## For Research Use Only