

A Reliable Research Partner in Life Science and Medicine

PE Anti-Mouse CD45R (B220) Antibody[RA3-6B2]

Catalog Number: AN00428UD

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Host Rat

Isotype Rat IgG2a, κ
Clone No. RA3-6B2

Isotype Control PE Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833D]

Conjugation PE

Conjugation Information PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green

(561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42

nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Data



Staining of C57BL/6 murine splenocytes cells with PE Anti-Mouse CD45R (B220) Antibody[RA3-6B2] (filled gray histogram) or PE Rat IgG2a, κ Isotype Control (empty black histogram). Total viable cells were used for analysis.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping lce bag

Antigen Information

 Alternate Names
 CD45R;B220

 Uniprot ID
 Q64224

 Gene ID
 19264;5788

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Background

CD45R, also known as B220, is an isoform of CD45. It is a member of the protein tyrosine phosphatase (PTP) family with a molecular weight of approximately 180-240 kD. CD45R is expressed on B cells (at all developmental stages from pro-B cells through mature B cells), activated B cells, and subsets of T and NK cells. CD45R (B220) is also expressed on a subset of abnormal T cells involved in the pathogenesis of systemic autoimmunity in MRL-FasIpr and MRL-Fasgld mice. It plays a critical role in TCR and BCR signaling. The primary ligands for CD45 are galectin-1, CD2, CD3, and CD4. CD45R is commonly used as a pan-B cell marker. CD19, however, may be more appropriate for B cell specificity.

Fax: 1-832-243-6017