Recombinant Rat LAG3/CD223 Protein (His Tag)

Catalog Number: PKSR030201

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Rat
Source	HEK293 Cells-derived Rat LAG3/CD223 protein Met1-Leu442, with an C-terminal His
Calculated MW	47.1 kDa
Observed MW	55 kDa
Accession	Q5BK54
Bio-activity	Not validated for activity
Properties	
Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 20 mM NaAc, 500 mM NaCl, 10 % Glycerol, 0.02 % tween
	20, pH 5.5
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	
	KDa M
	116
	66.2
	45.0
	35.0
	55.0
	25.0

> 90 % as determined by reducing SDS-PAGE.

18.4 14.4

Background

LAG3, also known as CD223 and Lymphocyte activation gene 3, belongs to immunoglobulin (Ig) superfamily. LAG3 contains 4 extracellular Ig-like domains. The LAG3 gene contains 8 exons. It is selectively expressed in activated T and NK cells. LAG3 has a negative regulatory function in T cells. It also acts as as a new marker of T cell induced B cell activation. As a soluble molecule, LAG3 activates antigen-presenting cells through MHC class II signalling, leading to increased antigen-specific T-cell responses in vivo.

Web:www.elabscience.com