



A Reliable Research Partner in Life Science and Medicine

Elab Fluor® Violet 450 Anti-Human CD64 Antibody[10.1]

Catalog Number: E-AB-F1082Q

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human Host Mouse

Isotype Mouse IgG1, κ

Clone No. 10.1

Isotype Control Elab Fluor® Violet 450 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-

F09792Q]

Conjugation Elab Fluor[®] Violet 450

Conjugation Information Elab Fluor[®] Violet 450 is designed to be excited by the violet laser (405 nm) and

detected using an optical filter centered near 450 nm (e.g., a 450/45 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD64;CD64A/B/C;FCGR1A/B/C;Fc fragment of lgG high affinity la/b/c receptor;Fc

gamma RI;IGFR 1

 Uniprot ID
 P12314

 Gene ID
 2209

Background CD64 is a 72 kD single chain type I glycoprotein also known as FcγRI and FcR I. CD64

Web: www.elabscience.cn

is a member of the immunoglobulin superfamily and is expressed on monocytes/macrophages, dendritic cells, and activated granulocytes. The expression can be upregulated by IFN-γ stimulation. CD64 binds IgG immune complex. It plays a role in antigen capture, phagocytosis of IgG/antigen complexes, and antibody-dependent

cellular cytotoxicity (ADCC).