

## APC Anti-Human IFN-γ Antibody[B27]

**Catalog Number:** E-AB-F1196E

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Isotype</b>	Mouse IgG1, κ
<b>Clone No.</b>	B27
<b>Isotype Control</b>	APC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792E]
<b>Conjugation</b>	APC
<b>Conjugation Information</b>	APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

### Applications

### Recommended usage

<b>FCM</b>	Each lot of this antibody is quality control tested by flow cytometric analysis. <b>The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).</b> Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
------------	---

### Data



HEK293T cells transiently transfected with pcDNA3.1 plasmid encoding Human IFN-γ gene are stained with APC Anti-Human IFN-γ Antibody (filled gray histogram) or APC Mouse IgG1, κ Isotype Control (empty black histogram).

### Preparation & Storage

<b>Storage</b>	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
<b>Shipping</b>	Ice bag

### Antigen Information

<b>Alternate Names</b>	IFN-gamma;IFNG;IFNγ;Interferon gamma
<b>Uniprot ID</b>	P01579
<b>Gene ID</b>	3458

### For Research Use Only

## Background

Interferon- $\gamma$  is a potent multifunctional cytokine which is secreted primarily by activated NK cells and T cells. Originally characterized based on anti-viral activities, IFN- $\gamma$  also exerts anti-proliferative, immunoregulatory, and proinflammatory activities. IFN- $\gamma$  can upregulate MHC class I and II antigen expression by antigen-presenting cells. The B27 antibody reacts with the human interferon- $\gamma$ . The B27 antibody can neutralize the bioactivity of natural or recombinant IFN- $\gamma$ .

## For Research Use Only