

A Reliable Research Partner in Life Science and Medicine

# APC Anti-Mouse CD206/MMR Antibody[C068C2]

Catalog Number: E-AB-F1135UE

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Mouse Rat Host

Isotype Rat IgG2a, ĸ Clone No. C068C2

APC Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833E] Isotype Control

Conjugation

**Conjugation Information** APC is designed to be excited by the Red (627-640 nm) laser and detected using an

optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).

Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer. Storage Buffer


#### **Applications** Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. Please **FCM** 

> check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 μg/10<sup>6</sup> cells

in 100 µL volume].

#### Data



C57BL/6 murine abdominal macrophages elicited by starch broth are stained with FITC Anti-Mouse F4/80 Antibody and APC Anti-Mouse CD206 Antibody (Left). Abdominal macrophages are stained with FITC Anti-Mouse F4/80 Antibody and APC Rat IgG2a, κ Isotype Control (Right).

#### **Preparation & Storage**

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

#### **Antigen Information**

**Alternate Names** MMR;MR;MRC1;macrophage mannose receptor;mannose receptor

**Uniprot ID** Q61830 Gene ID 17533

## For Research Use Only

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Toll-free: 1-888-852-8623

Email:techsupport@elabscience.com Web:www.elabscience.com

#### Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

## Background

Elabscience®

CD206, also known as mannose receptor (MR), is a 175 kD type I membrane protein. It is a pattern recognition receptor (PRR) belonging to the C-type lectin superfamily. MR is expressed on macrophages, dendritic cells, Langerhans cells, and hepatic or lymphatic endothelial cells. MR recognizes a range of microbial carbohydrates bearing mannose, fucose, or N-acetyl glucosamine through its C-type lectin-like carbohydrate recognition domains, sulfated carbohydrate antigens through its cysteine-rich domain, and collagens through its fibronectin type II domain. MR mediates endocytosis and phagocytosis as well as activation of macrophages and antigen presentation. It plays an important role in host defense and provides a link between innate and adaptive immunity. Recently, MR on lymphatic endothelial cells was found to be involved in leukocyte trafficking and a contributor to the metastatic behavior of cancer cells. It suggests that MR may be a potential target in controlling inflammation and cancer metastasis by targeting the lymphatic vasculature.

Fax: 1-832-243-6017