Recombinant Human BMP-2 Protein(Sumo Tag)

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Catalog Number: PDEH100503



Description Species Human Source E.coli-derived Human BMP-2 protein Gln283-Arg396, with an N-terminal Sumo Mol Mass 25.4 kDa P12643 Accession **Bio-activity** Not validated for activity **Properties** Purity >90% as determined by reducing SDS-PAGE. Endotoxin < 10 EU/mg of the protein as determined by the LAL method Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 Storage °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at $< -20^{\circ}$ C for 3 months. Shipping This product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Formulation Mannitol. Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data

kDa	м	R
80	-	
60	-	
40	-	
30		-
20	Η.	

SDS-PAGE analysis of Human BMP-2 proteins, 2 μg/lane of Recombinant Human BMP-2 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 25.4 KD

Background

Bone morphogenetic protein-2 (BMP-2) is a member of the transforming growth factor-beta (TGFB) superfamily. BMP2 is synthesized as a 60 kDa precursor that is processed in the secretory pathway to a small 18 kDa monomer, 2 monomers then associate to form the active 30 kDa homodimer, which binds to its receptor. There is also a 40-45 kDa form of BMP2, as an amino-terminal propeptide. BMP2 can induce bone formation and regeneration during early embryonic development. It is involved in the hedgehog pathway, TGF beta signaling pathway, and cytokine-cytokine receptor interaction.

For Research Use Only