

## Purified Anti-Human CD147 Antibody[HIM6]

catalog number: E-AB-F1056A

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


|                    |  |
|--------------------|--|
| <b>Reactivity</b>  | Human  |
| <b>Host</b>        | Mouse  |
| <b>Isotype</b>     | Mouse IgG1, $\kappa$   |
| <b>Clone</b>       | HIM6   |
| <b>Conjugation</b> | Unconjugated   |
| <b>Buffer</b>      | Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling. |

### Applications

### Recommended Dilution

|            |   |
|------------|---|
| <b>FCM</b> | 2 $\mu\text{g/mL}$ ( $1 \times 10^5$ - $5 \times 10^5$ cells) |
|------------|---|

### Data



Human peripheral blood lymphocytes were stained with 0.2  $\mu\text{g}$  Purified Anti-Human CD147 Antibody[HIM6] (Right) and 0.2  $\mu\text{g}$  Mouse IgG1,  $\kappa$  Isotype Control (Left), followed by FITC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

### Preparation & Storage

|                 |  |
|-----------------|--|
| <b>Storage</b>  | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. |
| <b>Shipping</b> | Ice bag  |

### Background

CD147, also known as neurothelin or basigin, is a member of the Ig superfamily. It is a 55-65 kD type I transmembrane glycoprotein which is primarily expressed on leukocytes, erythrocytes, platelets, and endothelial cells. CD147 is reported to have a function during embryonal brain development and/or play a role in integrin-mediated adhesion in brain endothelia.