

## Recombinant Human MMP19 Protein (His Tag)

**Catalog Number:** PDEH101035

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	E.coli-derived Human MMP19 protein Leu101-Arg450, with an N-terminal His & C-terminal His
<b>Calculated MW</b>	38.4 kDa
<b>Observed MW</b>	41 kDa
<b>Accession</b>	Q99542
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 90% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 10 EU/mg of the protein as determined by the LAL method
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



SDS-PAGE analysis of Human MMP19 proteins, 2 µg/lane of Recombinant Human MMP19 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 41 kDa.

### Background

MMP-19 (Matrix metalloprotease 19, also MMP-18 and MMP RASI) is a 55-59 kDa member of the peptidase M10A family of enzymes. It is widely expressed, being secreted by stratum basale keratinocytes, smooth muscle cells, epiphyseal cartilage chondrocytes and monocytes/macrophages. MMP-19 has multiple substrates, including components of the basement membrane (type IV collagen, laminin, nidogen), fibronectin, aggrecan plus COMP, and IGFBP3, this latter cleavage resulting in the release of active IGF-I. Studies involving MMP-19 demonstrate an antiangiogenic function. This is attributable to the processing of plasminogen, generating angiostatin-like molecules, and the creation of an environment that promotes the ECM retention of soluble VEGF.

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017