## Recombinant Mouse Osteonectin/SPARC Protein (His Tag)

Catalog Number: PKSM041280



Description		
Species	Mouse	
Mol Mass	33.6 kDa	
Accession	P07214	
Bio-activity	Not validated for activity	
Properties		
Purity	> 95 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80	
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of	
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.	
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants	
	before lyophilization.	
	Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	

Note: Centrifuge before opening to ensure complete recovery of vial contents.

kDa	MK	R
120 90 60		
40	-	
30		
20		
14		

## Background

Data

Secreted protein acidic and rich in cysteine (SPARC,BM-40) protein belongs to the family of secreted matricellular proteins with similar domain structure. Mouse SPARC protein involved an N-terminal acidic region that binds calcium, a follistatin domain containing Kazal-like sequences, and a C-terminal extracellular calcium (EC) binding domain with two EF-hand motifs. SPARC is produced by fibroblasts, capillary endothelial cells, platelets, and macrophages, especially in areas of tissue morphogenesis and remodeling. It appears to regulate cell growth through interactions with the extracellular matrix and cytokines. SPARC is expressed at high levels in tissues undergoing morphogenesis, remodeling and wound repair. The activity of SPARC is to modulate cell-cell and cell-matrix interactions, and its de-adhesive and growth inhibitory properties in non-transformed cells have led to studies to assess its role in cancer.

## For Research Use Only