

## M199

Cat. No.: PM150610

Size: 500mL

### General Information

<b>Product Form</b>	Liquid
<b>Concentration</b>	1 ×
<b>pH</b>	7.0-7.6
<b>D-Glucose</b>	1000 mg/L
<b>HEPES</b>	Negative
<b>L-Glutamine</b>	100 mg/L
<b>NaHCO<sub>3</sub></b>	2200 mg/L
<b>Phenol red</b>	20 mg/L
<b>Balanced salt</b>	Earle's salt
<b>Storage</b>	2-8°C, Shading Light
<b>Shipping</b>	Room Temperature
<b>Expiration date</b>	24 months

### Background

M199 (Medium 199) originally used in nutritional studies of chicken embryo fibroblasts, it has been widely used in the culture of various animal cells, including some non-mammalian cells. M199 is especially suitable for the culture of untransformed cells, rat pancreatic epithelial cells and mouse lens tissues, also used in virology and vaccine production. Compared with other basic medium, M199 contains unique ingredients, including adenine, adenosine, hypoxanthine, thymine, and other vitamins. There are two kinds of equilibrium salt (earle's salt and hank's salt) in M199. Earle's salt is often used in CO<sub>2</sub> environment, hank's salt is used in non-CO<sub>2</sub> environment.

This product contains many kinds of amino acids, vitamins, inorganic salts and other ingredients for cell culture, but does not contain protein, lipids or any growth factors. Therefore, the product should be used with serum or serum-free additives.

### Guidelines for use

Procell's cell culture media undergoes strict quality control to ensure sterility, but may get contaminated during use. Follow these guidelines for sterile handling to avoid contamination.

1. Always wipe your gloved hands and work area with 70% ethanol.
2. Wipe the outside of the containers, flasks, plates, and dishes with 70% ethanol before placing them in the cell culture hood.
3. Use sterile pipette tips and pipettes to work with liquids, and use each pipette tip only once to avoid crosscontamination. Do not unwrap sterile pipettes until they are ready to be used. Keep pipettes and tips within the clean work area.
4. Do not talk while performing sterile procedures and perform your cell culture as efficiently and carefully as possible to minimize contamination.

## Quality control

Standard evaluations for cell culture media are pH, osmolality, endotoxins and sterility testing for liquid products, cell growth experiments.

## Notes

1. This product is only used for scientific research or further research, not for diagnosis and treatment.
2. This product is sterilized by 0.1 µm filtration.
3. It is necessary to pay attention to the aseptic operation and avoid the contamination.

Procell system®

Procell system®

Procell system®

Procell system®