

## AF/LE Purified Anti-Mouse CD119 Antibody[GR-20]

catalog number: E-AB-F11150

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Mouse
<b>Immunogen</b>	Recombinant Mouse CD119 protein
<b>Host</b>	Rat
<b>Isotype</b>	Rat IgG2a, κ
<b>Clone</b>	GR-20
<b>Purification</b>	>98%, Protein A/G purified
<b>Buffer</b>	Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method.

### Applications

### Recommended Dilution

<b>FCM</b>	2 µg/mL(0.5×10 <sup>6</sup> -1×10 <sup>6</sup> cells)
------------	---

### Data



C57/BL6 Mouse splenocytes were stained with 0.2 µg AF/LE Purified Anti-Mouse CD119 Antibody[GR-20](Right) and 0.2 µg Rat IgG2a, κ Isotype Control (Left), followed by APC-conjugated Goat Anti-Rat IgG Secondary Antibody.

### Preparation & Storage

<b>Storage</b>	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions.
<b>Shipping</b>	Ice bag

### Background

CDw119 is a 90 kD immunoglobulin superfamily member, also known as IFN-γRα chain. It is a class II cytokine receptor family member that serves as a IFN-γ-binding chain associated with the IFN-γβ chain also known as AF-1. In addition to ligand binding, CDw119 participates in ligand trafficking. CDw119 is expressed on T and B cells, NK cells, fibroblasts, endothelial, and epithelial cells. Binding of IFN-γ induces receptor dimerization, internalization, Jak1 and Jak2 protein kinase activation and, ultimately, STAT1 activation. IFN-γ initiates and regulates a variety of immune responses.

### For Research Use Only