ARMC8 Polyclonal Antibody

catalog number: E-AB-18698



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse
Immunogen	Fusion protein of human ARMC8
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
IHC	1:40-1:200
Data	
esophagus cancer tissue u at diluti	of paraffin-embedded Human Immunohistochemistry of paraffin-embedded Human sing ARMC8 Polyclonal Antibody thyroid cancer tissue using ARMC8 Polyclonal Antibody at on of 1:40(×200) dilution of 1:40(×200)
Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the
	temperature recommended.
Background	
The armadillo (ARM) rep	eat family of proteins are related to the Drosophila melanogaster armadillo protein, a protein

essential for wingless signal transduction. ARM proteins are related to the Diosophila metalogaster annaulio protein, a protein essential for wingless signal transduction. ARM proteins are involved in a variety of processes such as cell migration, cell proliferation, tissue maintenance and tumorigenesis, and they also function in signal transduction and the maintenance of overall cell structure. ARMC8 (armadillo repeat containing 8), also known as S863-2, is a 673 amino acid protein that contains 14 ARM repeats, suggesting a role in signal transduction events throughout the cell. Six isoforms of ARMC8 are expressed due to alternative splicing events. The gene encoding ARMC8 maps to human chromosome 3, which houses over 1,100 genes, including a chemokine receptor (CKR) gene cluster and a variety of human cancerrelated gene loci.

For Research Use Only