## SYT4 Polyclonal Antibody

catalog number: E-AB-14373



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Reactivity	Human;Mouse;Rat		
Immunogen	Recombinant protein of human SYT4		
Host	Rabbit		
Isotype	IgG		
Purification	Affinity purification		
Conjugation	Unconjugated		
buffer	Phosphate buffered solu	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.	
Applications	Recommended Dilution		
WB	1:500-1:2000	1:500-1:2000	
IHC	1:25-1:100		
Data			
60-1 137- 7- 5- 5-	-		
Western Blot analysis of Hela	•	Immunohistochemistry of paraffin-embedded Human colon	
Polyclonal Antibody at dilution of 1:700		cancer using SYT4 Polyclonal Antibody at dilution of 1:60	
Calculated-MV:48 kDa			
Preparation & Storage			
Storage	Store at -20°C Valid for 12	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.	
Shipping	The product is shipped w	The product is shipped with ice pack, upon receipt, store it immediately at the	
	temperature recommended.		
Background			

Synaptotagmins are a large gene family of synaptic vesicle type III integral membrane proteins that function as regulators of both exocytosis and endocytosis and are involved in neurotransmitter secretion from small secretory vesicles. Calcium binds to Synaptotagmin I which triggers neurotransmitter release at the synapse. Synaptotagmin II is phosphorylated by WNK1 in a process that regulates calcium-dependent interactions. Synaptotagmin IV is expressed in calcium-dependent exocytosis of secretory vesicles in endocrine cells and neurons. Synaptotagmin IV is expressed in neuronal tissues, and has the highest mRNA levels in the hippocampus. The proximity of the Synaptotagmin IV gene to markers of several psychiatric disorders suggest an involvement of synaptotagmin IV in human disease. Synaptotagmin V is a dense-core vesicle-specific protein that regulates a specific type of calcium-regulated secretion. Synaptotagmin VI interacts with adaptor protein-2 in a calcium-independent manner. Synaptotagmin VII is widely expressed in non-neuronal tissues.

## For Research Use Only