

GM-CSF/CSF2/CSF, Human, Recombinant

Cat. No. : PCK035

General Information

Synonyms	colony stimulating factor 2;CSF2;CSF
Species	Human
Expression host	E.coli
Sequence	APARSPSPSTQPWEHVNAIQEARRLLNLSRDTAEMNETVEVISEMFDLQEPTCLQTRLELYKQ GLRGLSTKLKGPLTMMASHYKQHCPTPETSCATQIITFESFKENLKDFLLVIPFDCWEPVQE with polyhistidine tag at the N-terminus.
Accession	P04141.1
Tag	His-tag at the N-terminus
Mol mass	15.4 kDa
Expiration date	12 months
Bio activity	Measure by its ability to induce TF-1 cells proliferation.The ED50 for this effect is < 80 pg/mL. The specific activity of recombinant human GM-CSF is approximately > 1 × 10 ⁷ IU/mg.

Product feature

Purity	> 98% as determined by SDS-PAGE. Ni-NTA chromatography.
Endotoxin (EU/μg)	< 0.1
Storage	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
Shipping	Ice bag
Formulation	The protein was lyophilized from a 0.2 μm filtered solution containing 1 × PBS, pH 8.0.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

Background

Granulocyte-macrophage colony-stimulating factor (GM-CSF), also known as colony-stimulating factor 2 (CSF2), is a monomeric glycoprotein secreted by macrophages, T cells, mast cells, natural killer cells, endothelial cells and fibroblasts that functions as a cytokine. GM-CSF also plays a role in embryonic development by functioning as an embryokine produced by reproductive tract.