# **Elabscience**®

### FITC Anti-Mouse H-2 Antibody[M1/42]


#### Catalog Number: E-AB-F1216UC

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Mouse
Host	Rat
lsotype	Rat lgG2a, κ
Clone No.	M1/42
Isotype Control	FITC Rat IgG2a, к Isotype Control[2A3] [Product E-AB-F09833C]
Conjugation	FITC
Conjugation Information	FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical filter centered near 530 nm (e.g., a 525/40 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 µg/10 <sup>6</sup> cells

reagent to obtain optimal results [The recommended concentration is 0.1-1  $\mu$ g/10<sup>o</sup> ce in 100  $\mu$ L volume].

Data



C57BL/6 murine splenocytes are stained with FITC Anti-Mouse H-2 Antibody (filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.

Keep as concentrated solution.
This product can be stored at 2-8°C for 12 months. Please protected from prolonged
exposure to light and do not freeze.
Ice bag
MHC I;Mouse major histocompatibility complex (MHC) H-2
P06345
111364

### For Research Use Only

## **Elabscience**®

Background

The M1/42 antibody reacts with the H-2 MHC class I alloantigens expressed on nucleated cells from mice of the a, b, d, j, k, s, and u haplotypes (Stallcup, KC et al, 1981). MHC class I is involved in antigen presentation to T cells expressing CD3/TCR and CD8 proteins.

#### For Research Use Only