

## Recombinant Arginase-1 Monoclonal Antibody

catalog number: **AN300873L**

**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*

### Description


|                     |                                                                 |
|---------------------|-----------------------------------------------------------------|
| <b>Reactivity</b>   | Human;Mouse;Rat                                                 |
| <b>Immunogen</b>    | Recombinant Human Arginase-1 protein                            |
| <b>Host</b>         | Rabbit                                                          |
| <b>Isotype</b>      | Rabbit IgG                                                      |
| <b>Clone</b>        | C110497                                                         |
| <b>Purification</b> | Protein A                                                       |
| <b>Buffer</b>       | PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant. |

### Applications

### Recommended Dilution

|              |                |
|--------------|----------------|
| <b>IHC</b>   | 1:200-1:1000   |
| <b>WB</b>    | 1:2000-1:10000 |
| <b>IF</b>    | 1:200-1:1000   |
| <b>ELISA</b> | 1:5000-1:20000 |
| <b>IP</b>    | 1:50-1:200     |


### Data



Western Blot with Recombinant Arginase-1 Monoclonal Antibody at dilution of 1:1000 dilution. Lane A: huh-7 cells.

**Observed-MW:35 kDa**

**Calculated-MW:35 kDa**



Immunohistochemistry of paraffin-embedded rat brain using Recombinant Arginase-1 Monoclonal Antibody at dilution of 1:200.

### Preparation & Storage

|                 |                                                                 |
|-----------------|-----------------------------------------------------------------|
| <b>Storage</b>  | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| <b>Shipping</b> | Ice bag                                                         |

### Background

Arginase catalyzes the hydrolysis of arginine to ornithine and urea. At least two isoforms of mammalian arginase exist (types I and II) which differ in their tissue distribution, subcellular localization, immunologic crossreactivity and physiologic function. The type I isoform encoded by this gene, is a cytosolic enzyme and expressed predominantly in the liver as a component of the urea cycle. Inherited deficiency of this enzyme results in argininemia, an autosomal recessive disorder characterized by hyperammonemia. Two transcript variants encoding different isoforms have been found for this gene.

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.4