# **Elabscience**®


### PE Anti-Mouse CD366/Tim-3 Antibody[RMT3-23]

#### Catalog Number: E-AB-F1192D

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Mouse
Host	Rat
lsotype	Rat IgG2a, ĸ
Clone No.	RMT3-23
Isotype Control	PE Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832D]
Conjugation	PE
<b>Conjugation Information</b>	PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green
	(561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42
	nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount
	of the reagent is suggested to be used 5 $\mu$ L of antibody per test (million cells in 100
	μL staining volume or per 100 μL of whole blood). Please check your vial before the
	experiment. Since applications vary, the appropriate dilutions must be determined for
	individual use.

Data



C57BL/6 murine splenocytes are stained with APC Anti-Mouse CD4 Antibody and PE Anti-Mouse CD366/Tim-3 Antibody (Left). Splenocytes are stained with APC Anti-Mouse CD4 Antibody and PE Rat IgG2a,  $\kappa$  Isotype Control (Right).

Preparation & Storage	e
Storage	Keep as concentrated solution.
	This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	TIMD3;HAVcr-2;TIM3;TIMD-3
Uniprot ID	Q8VIM0

### For Research Use Only

## **Elabscience**®

Gene ID Background

#### 84868

CD366 (Tim-3) is a transmembrane protein also known as T cell immunoglobulin and mucin domain containing protein-3. Tim-3 is expressed at high levels on activated T cells (preferentially on Th1 cells, monocytes/macrophages, and dendritic cells). Tim-3 has also been shown to exist as a soluble protein. Cells expressing Tim-3 are present at high levels in the CNS of animals at the onset of experimental autoimmune encephalomyelitis (EAE), a disease mediated by lymphocytes secreting Th1-like cytokines. Tim-3 has been proposed to inhibit Th1-mediated immune responses and promote immunological tolerance.