

A Reliable Research Partner in Life Science and Medicine

PerCP Anti-Human HLA-DR Antibody[L243]

Catalog Number: E-AB-F1111F

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human Host Mouse

Isotype Mouse IgG2a, κ

Clone No. L243

Isotype Control PerCP Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802F]

Conjugation PerCP

Conjugation Information PerCP is designed to be excited by the blue laser (488 nm) and detected using an optical

filter centered near 675 nm (e.g., a 690/50 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein

protectant.


Applications

Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Human peripheral blood lymphocytes are stained with PerCP Anti-Human HLA-DR Antibody (filled gray histogram). Unstained lymphocytes (empty black histogram) are used as control.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping lce bag

Antigen Information

Alternate Names DRA/DRB1;HLA class II histocompatibility antigen DR alpha/ DRB1-15 beta chain;HLA-

DRA1/DRB1;MHC class II antigen DRA

Uniprot ID P01903;P01911

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web:www.elabscience.com
 Email:techsupport@elabscience.com

Elabscience Bionovation Inc.

Elabscience®

A Reliable Research Partner in Life Science and Medicine

Gene ID

Background

3122,3123

HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36 kD α (heavy) chain and a 27 kD β (light) chain. It is expressed on B cells, activated T cells, monocytes/macrophages, dendritic cells, and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4+ T cells.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web:www.elabscience.com
 Email:techsupport@elabscience.com