

Recombinant Human CD69 Protein (aa 64-199, His Tag)



Catalog Number:PKSH033694

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Early activation antigen CD69;Activation inducer molecule;AIM;BL-AC/P26;C-type lectin domain family 2 member C;EA1;Early T-cell activation antigen p60;GP32/28;Leukocyte surface antigen Leu-23;MLR-3;CD69;CLEC2C
Species	Human
Expression Host	HEK293 Cells
Sequence	Gly64-Lys199
Accession	Q07108
Calculated Molecular Weight	16.9 kDa
Observed molecular weight	18-28 kDa
Tag	N-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Human Early Activation Antigen CD69 (CD69) is a type 2 transmembrane glycoprotein in the C-type lectin family. It plays roles in immune cell trafficking, inflammation, T cell memory, and humoral immune responses. CD69 is expressed on the cell surface as an approximately 60 kDa disulfide-linked homodimer. It is found on CD4+ T cells, CD8+ T cells, NK cells, NKT cells, gamma delta cells dendritic cells (DC) and is up-regulated on activated T cells and DC. Ligation of CD69 on DC induces IL2 production, leading to T cell proliferation. CD69 is important for the homing of CD4+ T cells and plasmablasts to the bone marrow but inhibits the migration of dermal DC to draining lymph nodes. It supports the

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human CD69 Protein (aa 64-199, His Tag)



Catalog Number: PKSH033694

expression of multiple chemokines and chemokine receptors but suppresses the expression of others. It associates with and negatively regulates S1P1 expression on DC and CD4+ T cells, resulting in a decreased chemotactic response to S1P. The direct interaction of CD69 with Galectin-1 contributes to the ability of CD69 to limit Th17 mediated inflammation while supporting the differentiation of regulatory T cells.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017