



A Reliable Research Partner in Life Science and Medicine

Recombinant Phospho-PLCy1 (Tyr783) Monoclonal Antibody

catalog number: AN301390L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat

Immunogen A synthetic peptide corresponding to residues around (Tyr783) of Human Phospho-

PLC_v1

 Host
 Rabbit

 Isotype
 IgG,κ

 Clone
 B1157


 Purification
 Protein A

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications Recommended Dilution

WB 1:2000-1:10000

Data



Western Blot with Recombinant Phospho-PLCγ1 (Tyr783) Monoclonal Antibody at dilution of 1:1000 dilution. Lane A:

Hela.

Observed-MW:149 kDa Calculated-MW:149 kDa

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

Background

The protein encoded by this gene catalyzes the formation of inositol 1,4,5-trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of receptor-mediated tyrosine kinase activators. For example, when activated by SRC, the encoded protein causes the Ras guanine nucleotide exchange factor RasGRP1 to translocate to the Golgi, where it activates Ras. Also, this protein has been shown to be a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)-activated tyrosine kinase. Two transcript variants encoding different isoforms have been found for this gene.

For Research Use Only