Recombinant Human Decorin Protein (Human Cells, His Tag)

Catalog Number: PKSH033700



Note: Centrifuge before opening to ensure complete recovery of vial contents.

			•	7.0	
	es	OF	TIM	tio	m
JU.	\mathbf{c}	UЦ	ш	ww	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,

 Species
 Human

 Mol_Mass
 38.8 kDa

 Accession
 P07585

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Decorin is a secreted chondroitin/dermatan sulfate proteoglycan in the family of small leucine-rich proteoglycans (SLRP s). SLRP family members are characterized by N-terminal and C-terminal cysteine-rich regions which flank the central region containing 10-12 tandem leucine-rich repeats (LRR). The human Decorin cDNA encodes a 359 amino acid (aa) precursor that includes a 16 aa signal sequence and a 14 aa propeptide. Alternate splicing of human Decorin generates five isoforms with variable length deletions. Decorin is an N-glycosylated protein that also carries a variablysized hybrid chondroitin/dermatan sulfate chain at Ser34. Decorin regulates assembly of the extracellular collagen matrix and the bioactivity of the matrix associated growth factors FGF2;GDF8/Myostatin; TGFβ;and WISP1. It also binds and activates EGF R; ErbB4; and IGFI-R. In vivo; Decorin promotes myoblast differentiation; supports angiogenesis; and inhibits tumor progression.

For Research Use Only