

Purified Anti-Human CD9 Antibody[HI9a]

catalog number: E-AB-F1086A

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


| | |
|---------------------|--|
| Reactivity | Human |
| Immunogen | Recombinant Human CD9 protein |
| Host | Mouse |
| Isotype | Mouse IgG1, κ |
| Clone | HI9a |
| Purification | >98%, Protein A/G purified |
| Buffer | Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling. |

Applications

Recommended Dilution

| | |
|------------|---|
| FCM | 2 μ g/mL (1×10^5 - 5×10^5 cells) |
|------------|---|

Data



Human peripheral blood platelets were stained with 0.2 μ g Purified Anti-Human CD9 Antibody[HI9a] (Right) and 0.2 μ g Mouse IgG1, κ Isotype Control (Left), followed by Alexa Fluor® 647-conjugated Goat Anti-Mouse IgG Secondary Antibody.

Preparation & Storage

| | |
|-----------------|--|
| Storage | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. |
| Shipping | Ice bag |

Background

CD9 is a 24 kD type III transmembrane protein also known as tetraspanin, MRP-1 and DRAP-24. It is a member of the tetraspan family (spanning the membrane four times) found on platelets, B cell progenitors, activated lymphocytes, granulocytes, endothelial cells and epithelial cells. CD9 induces adhesion, platelet aggregation, and B cell development. CD9 has been shown to associate with CD63, CD81, CD82, and CD36 and to bind to β 1 integrins.

For Research Use Only