



A Reliable Research Partner in Life Science and Medicine

Biotin Anti-Mouse CD127/IL-7RA Antibody[A7R34]

Catalog Number: E-AB-F1023B

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Host Rat

Isotype Rat IgG2a, κ **Clone No.** A7R34

Isotype Control Biotin Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833B]

Conjugation Biotin

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. For flow

cytometric staining, the suggested use of this reagent is \leq 1.0 μ g per 10⁶ cells in 100 μ L volume or 100 μ L of whole blood. It is recommended that the reagent be titrated for

optimal performance for each application.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD127;IL-7 receptor subunit alpha;IL-7R subunit alpha;IL-7R-alpha;IL-7RA;II7r;

Interleukin-7 receptor subunit alpha

 Uniprot ID
 P16872

 Gene ID
 16197

Background CD127 is a 60-90 kD type I transmembrane glycoprotein also known as IL-7 receptor α

chain or IL-7R α . It forms a heterodimer with the common γ chain (γ c or CD132) which is

shared with the receptors for IL-2, IL-4, IL-9, IL-13, IL-15, and IL-21. CD127 is expressed on immature B cells through early pre-B stage, thymocytes (except CD4/CD8 double positive thymocytes), peripheral T cells, and bone marrow stromal cells. CD127 has been reported to be an useful marker for identifying memory and effector T cells. The ligation of IL-7 with its receptor is important for stimulation of mature and immature T cells as well as immature B cells proliferation and development.

For Research Use Only