

Recombinant Human CD16a/FCGR3A Protein (His Tag)



Catalog Number:PKSH032420

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Low Affinity Immunoglobulin Gamma Fc Region Receptor III-A;CD16a Antigen;Fc-Gamma RIII-Alpha;Fc-Gamma RIII;Fc-gamma RIIIa;FcRIII;FcRIIIa;FcR-10;IgG Fc Receptor III-2;CD16a;FCGR3A;CD16A;FCG3;FCGR3;IGFR3;CD16;CD16A
Species	Human
Expression Host	HEK293 Cells
Sequence	Gly17-Gln208
Accession	P08637
Calculated Molecular Weight	22.7 kDa
Observed molecular weight	35-50 kDa
Tag	C-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Receptors for the Fc region of immunoglobulin G (Fc γ R) are divided into three classes and Fc γ RIII is a multifunctional; low/intermediate affinity receptor. In humans; Fc γ RIII is expressed as two distinct forms (Fc γ RIIIA and Fc γ RIIB) that are encoded by two different but highly homologous genes in a cell type-specific manner. Fc γ RIIB is a low-affinity; GPI-linked receptor expressed by neutrophils and eosinophils; whereas Fc γ RIIIA is an intermediate affinity polypeptide-anchored transmembrane glycoprotein expressed by a subset of T lymphocytes; natural killer (NK) cells; monocytes; and

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human CD16a/FCGR3A Protein (His Tag)



Catalog Number:PKSH032420

macrophages. The Fc γ RIIIA receptor is involved in phagocytosis; secretion of enzymes; inflammatory mediators; antibody-dependent cellular cytotoxicity (ADCC); mast cell degranulation; and clearance of immune complexes.

Fc γ RIIIA has an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain and delivers an activation signal in the immune responses. Aberrant expression or mutations in this gene is implicated in susceptibility to recurrent viral infections; systemic lupus erythematosus; and alloimmune neonatal neutropenia. In humans; it is a 50 -70 kD type I transmembrane activating receptor.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017