Recombinant Cynomolgus TIM-3/HAVCR2 Protein (Fc Tag)



Catalog Number: PKSQ050022

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms T cell immunoglobulin and mucin domain3;HAVCR2;Tim-3;TIM3

Species Cynomolgus macaques

Expression Host

Sequence

Ser22-Arg201

Accession

G7P6Q7

Calculated Molecular Weight

Observed molecular weight

Tag

HEK293 Cells

Ser22-Arg201

46.3 kDa

55-75 kDa

C-Fc

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin $< 1.0 \text{ EU per } \mu \text{g of the protein as determined by the LAL method.}$

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

T cell immunoglobulin and mucin domain 3 is a member of the TIM family of immune regulating molecules. Mature cynomolgus TIM3 consists of a 182 amino acid (aa)extracellular domain (ECD), a 21 aa transmembrane segment, and a 78 aa cytoplasmic tail. TIM3 is up-regulated on several populations of activated myeloid cells (macrophage, monocyte, dendritic cell, microglia, mast cell) and T cells (Th1, CD8+, NK, Treg). Its binding to Galectin9 induces a range of immunosuppressive functions which enhance immune tolerance and inhibit anti-tumor immunity. TIM3 ligation attenuates CD8+ and Th1 cell responses and promotes the activity of Treg and myeloid derived suppressor cells. TIM3 interactions with Galectin-9 can trigger immune stimulatory effects, such as the coactivation of NK cell cytotoxicity.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com