

A Reliable Research Partner in Life Science and Medicine

Purified Anti-Mouse CD38 Antibody[NIMR5]

catalog number: E-AB-F1193A

Note: Centrifuge before opening to ensure complete recovery of vial contents.

T .		4.
Descr	Tm	ION
DUSUI	T D	шош

Reactivity Mouse **Host** Rat


Buffer Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze

to completely remove the stabilizer prior to labeling.

Applications Recommended Dilution

FCM $2 \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$

Data



C57/BL6 Mouse splenocytes were stained with 0.2 μ g Purified Anti-Mouse CD38 Antibody[NIMR5] (Right) and 0.2 μ g Rat IgG2a, κ Isotype Control (Left), followed by

FITC-conjugated Goat Anti-Rat IgG Secondary Antibody.

Preparation & Storage

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping Ice bag

Background

CD38 is a 42 kD glycoprotein, also known as T10. It is an ADP-ribosyl hydrolase, expressed on B cells, NK cells, a subset of T cells, brain, muscle, and kidney. In mouse, CD38 expression is downregulated on germinal center B cells and plasma cells, whereas this is not the case for humans. By functioning as both a cyclase and a hydrolase, CD38 mediates lymphocyte activation, as well as adhesion and metabolism of cADPR and NAADP. CD31 is the ligand of CD38.